

Dosing, Administration, and Billing Guide

Indications

- ZERBAXA is indicated for the treatment of adult patients (18 years and older) with hospital-acquired bacterial pneumonia and ventilator-associated bacterial pneumonia (HABP/VABP), caused by the following susceptible Gram-negative microorganisms: Enterobacter cloacae, Escherichia coli, Haemophilus influenzae, Klebsiella oxytoca, Klebsiella pneumoniae, Proteus mirabilis, Pseudomonas aeruginosa, and Serratia marcescens.
- ZERBAXA is indicated for the treatment of adult and pediatric patients (birth to less than 18 years old) with complicated urinary tract infections (cUTI), including pyelonephritis, caused by the following susceptible Gram-negative microorganisms: Escherichia coli, Klebsiella pneumoniae, Proteus mirabilis, and Pseudomonas aeruginosa.
- ZERBAXA used in combination with metronidazole is indicated for the treatment of adult and pediatric patients (birth to less than 18 years old) with complicated intra-abdominal infections (cIAI) caused by the following susceptible Gram-negative and Gram-positive microorganisms: Enterobacter cloacae, Escherichia coli, Klebsiella oxytoca, Klebsiella pneumoniae, Proteus mirabilis, Pseudomonas aeruginosa, Bacteroides fragilis, Streptococcus anginosus, Streptococcus constellatus, and Streptococcus salivarius.

Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of ZERBAXA and other antibacterial drugs, ZERBAXA should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Important Safety Information

 Patients with renal impairment: Decreased efficacy of ZERBAXA has been observed in patients with baseline CrCl of 30 to ≤50 mL/min. In a clinical trial of adult patients, patients with cIAIs with CrCl >50 mL/min had a clinical cure rate of 85.2% when treated with ZERBAXA plus metronidazole vs 87.9% when treated with meropenem. In the same trial, patients with CrCl 30 to \leq 50 mL/min had a clinical cure rate of 47.8% when treated with ZERBAXA plus metronidazole vs 69.2% when treated with meropenem. A similar trend was also seen in the cUTI trial. Dose adjustment is required for adult patients with CrCl 50 mL/min or less. All doses of ZERBAXA are administered over 1 hour. Monitor CrCl at least daily in patients with changing renal function and adjust the dose of ZERBAXA accordingly.

Please see additional Important Safety Information on the following pages and accompanying Prescribing Information.

Dosage and Administration

Preparation of Solutions

How ZERBAXA Is Supplied

Contents

Dosage and Administration
Preparation of Solutions
How ZERBAXA [®] (ceftolozane and tazobactam) Is Supplied
Coding and Billing Information



•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		3	
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	(6	
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		8	
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		9)



Dosage and Administration

Recommended Dosage in Adult Patients

The recommended dosage of ZERBAXA® (ceftolozane and tazobactam) in adult patients (18 years and older) with creatinine clearance (CrCl) greater than 50 mL/min is 1.5 gram (g) (ceftolozane 1 g and tazobactam 0.5 g) for cIAI and cUTI and 3 grams (ceftolozane 2 g and tazobactam 1 g) for HABP/VABP administered every 8 hours by intravenous infusion over 1 hour. The duration of therapy should be guided by the severity and site of infection and the patient's clinical and bacteriological progress, as shown below.

Dosage of ZERBAXA by Infection in Adult Patients (18 Years and Older) With CrCl^a >50 mL/min

Infection	Dose	Frequency	Infusion Time	Duration of Treatment
Complicated intra-abdominal infections ^b	1.5 g	Every 8 hours	1 hour	4 to 14 days
Complicated urinary tract infections, including pyelonephritis	1.5 g	Every 8 hours	1 hour	7 days

Infection	Dose	Frequency	Infusion Time	Duration of Treatment
Hospital-acquired bacterial pneumonia and ventilator- associated bacterial pneumonia (HABP/VABP)	3 g	Every 8 hours	1 hour	8 to 14 days

IV, intravenous.

^aCrCl estimated using Cockcroft-Gault formula.

^bUsed in conjunction with metronidazole 500 mg intravenously every 8 hours.

Important Safety Information (continued)

 Hypersensitivity: ZERBAXA is contraindicated in patients with known serious hypersensitivity to the components of ZERBAXA (ceftolozane/ tazobactam), piperacillin/tazobactam, or other members of the beta-lactam class. Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving beta-lactam antibacterials. Before initiating therapy with ZERBAXA, make careful inquiry about previous hypersensitivity reactions to cephalosporins, penicillins, or other beta-lactams. If an anaphylactic reaction to ZERBAXA occurs, discontinue use and institute appropriate therapy.



Not actual size.



Preparation of Solutions

DOSAGE AND

ADMINISTRATION



Is Supplied

How

ZERBAXA

Dosage and Administration (continued)

Recommended Dosage in Pediatric Patients With cIAI or cUTI (Birth to Less Than 18 Years of Age)

The recommended dosage regimen of ZERBAXA in pediatric patients from birth to less than 18 years of age with cIAI and cUTI with an estimated glomerular filtration rate (eGFR) greater than 50 mL/min/1.73m² is described below. ZERBAXA is administered every 8 hours by intravenous infusion over 1 hour. The duration of treatment should be guided by the severity and site of infection and the patient's clinical and bacteriological progress as shown below. For the treatment of cIAI, metronidazole should be given concurrently.

There is insufficient information to recommend a dosage regimen for pediatric patients with HABP/VABP.

Dosage of ZERBAXA by Infection in Pediatric Patients (Birth to Less Than 18 Years of Age) With eGFR^a Greater Than 50 mL/min/1.73m²

Infection	Dose	Frequency	Infusion Time	Duration of Treatment
Complicated intra-abdominal infections ^b	30 mg/kg up to a maximum dose of 1.5 g°	Every 8 hours	1 hour	5 to 14 days
Complicated urinary tract infections including pyelonephritis	30 mg/kg up to a maximum dose of 1.5 g°	Every 8 hours	1 hour	7 to 14 days

eGFR, estimated glomerular filtration rate.

^aEstimated GFR using an age-appropriate equation for use in the pediatric population.

^bUsed in conjunction with metronidazole.

^oPediatric patients weighing greater than 50 kg should not exceed a maximum dose of 1.5 g.

Important Safety Information (continued)

- Clostridioides difficile-associated diarrhea (CDAD), ranging from mild diarrhea to fatal colitis, has been reported with nearly all systemic antibacterial agents, including ZERBAXA. Careful medical history is necessary because CDAD has been reported to occur more than 2 months after the administration of antibacterial agents. If CDAD is confirmed, antibacterial use not directed against C. difficile should be discontinued, if possible.
- Development of drug-resistant bacteria: Prescribing ZERBAXA in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.
- Adverse reactions in adult patients with HABP/VABP: The most common adverse reactions occurring in ≥5% of adult patients receiving ZERBAXA in the HABP/VABP trial were hepatic transaminase increased (11.9%), renal impairment/renal failure (8.9%), and diarrhea (6.4%).

Preparation of Solutions

DOSAGE AND

ADMINISTRATION



How ZERBAXA **Is Supplied**

Dosage and Administration (continued)

Dosage Adjustments in Adult Patients With Renal Impairment

Dose adjustment is required for adult patients (18 years and older) with CrCl 50 mL/min or less, as shown below. All doses of ZERBAXA[®] (ceftolozane and tazobactam) are administered over 1 hour.

For patients with changing renal function, monitor CrCl at least daily and adjust the dosage of ZERBAXA accordingly.

Dosage of ZERBAXA in Adult Patients (18 Years and Older) With Renal Impairment (CrCl ≤50 mL/min)

Estimated CrCl (mL/min)ª	Complicated Intra-abdominal Infections and Complicated Urinary Tract Infections, Including Pyelonephritis	Hospital-acquired Bacterial Pr associated Bacterial Pneumo
30 to 50	750 mg (500 mg and 250 mg) intravenously every 8 hours	1.5 g (1 g and 0.5 g) intravenou
15 to 29	375 mg (250 mg and 125 mg) intravenously every 8 hours	750 mg (500 mg and 250 mg)
End-stage renal disease (ESRD) on hemodialysis (HD)	A single loading dose of 750 mg (500 mg and 250 mg) followed by a 150 mg (100 mg and 50 mg) maintenance dose administered intravenously every 8 hours for the remainder of the treatment period (on hemodialysis days, administer the dose at the earliest possible time following completion of dialysis)	A single loading dose of 2.25 g (450 mg (300 mg and 150 mg) r intravenously every 8 hours for t period (on hemodialysis days, ac possible time following completi

CrCl, creatinine clearance.

^aCrCl estimated using Cockcroft-Gault formula.

Dosage Adjustments in Pediatric Patients With Renal Impairment

Dosage adjustment of ZERBAXA in pediatric patients (birth to less than 18 years of age) with eGFR 50 mL/min/1.73 m² or less has not been determined. ZERBAXA is not recommended in pediatric patients who have an eGFR 50 mL/min/1.73m² or less.

Important Safety Information (continued)

- Adverse reactions in adult patients with cIAI or cUTI: The most common adverse reactions occurring in ≥5% of adult patients receiving ZERBAXA in the cUTI and cIAI trials were headache (5.8%) in the cUTI trial, and nausea (7.9%), diarrhea (6.2%), and pyrexia (5.6%) in the cIAI trial.
- Adverse reactions in pediatric patients with cIAI or cUTI: The most common adverse reactions occurring in \geq 7% of pediatric patients receiving ZERBAXA in the cIAI trial were diarrhea (17%), thrombocytosis (16%), pyrexia (13%), abdominal pain (11%), vomiting (10%), increased aspartate aminotransferase (7%), and anemia (7%). The most common adverse reactions occurring in \geq 7% of pediatric patients receiving ZERBAXA in the cUTI trial were thrombocytosis (9%), leukopenia (8%), diarrhea (7%), and pyrexia (7%).

neumonia and Ventilatornia

usly every 8 hours

intravenously every 8 hours

1.5 g and 0.75 g) followed by a maintenance dose administered he remainder of the treatment dminister the dose at the earliest ion of dialysis)

DOSAGE AND ADMINISTRATION

Preparation of Solutions



How ZERBAXA **Is Supplied**

Preparation of Solutions

ZERBAXA® (ceftolozane and tazobactam) does not contain a bacteriostatic preservative. Aseptic technique must be followed in preparing the infusion solution.



Preparation of Doses

ZERBAXA® (ceftolozane and tazobactam) Dose	Volume to Withdraw From Reconstituted Vial(s)
3 g (2 g and 1 g)	Two vials of 11.4 mL each (entire contents from two vials)
2.25 g (1.5 g and 0.75 g)	11.4 mL from one vial (entire contents) and 5.7 mL from a se
1.5 g (1 g and 0.5 g)	11.4 mL (entire contents from one vial)
750 mg (500 mg and 250 mg)	5.7 mL
450 mg (300 mg and 150 mg)	3.5 mL
375 mg (250 mg and 125 mg)	2.9 mL
150 mg (100 mg and 50 mg)	1.2 mL

Inspect drug products visually for particulate matter and discoloration prior to use. ZERBAXA infusions range from clear, colorless solutions to solutions that are clear and slightly yellow. Variations in color within this range do not affect the potency of the product.

Important Safety Information (continued)

• Pediatric Use: There is insufficient information to recommend dosage adjustment for pediatric patients younger than 18 years of age with cIAI and cUTI with eGFR 50 mL/min/1.73m² or less. ZERBAXA is not recommended in pediatric patients who have an eGFR 50 mL/min/1.73m² or less. Pediatric patients born at term or pre-term may not have an eGFR of 50 mL/min/1.73m² or greater at birth or within the first few months of life.

econd vial



Dosage and Administration

PREPARATION **OF SOLUTIONS**

How ZERBAXA Is Supplied

Preparation of Solutions (continued)

Compatibility

Compatibility of ZERBAXA® (ceftolozane and tazobactam) with other drugs has not been established. ZERBAXA should not be mixed with other drugs or physically added to solutions containing other drugs.

Storage of Constituted Solutions

Upon constitution with sterile water for injection or 0.9% sodium chloride injection, reconstituted ZERBAXA solution may be held for 1 hour prior to transfer and dilution in a suitable infusion bag.

Following dilution of the solution with 0.9% sodium chloride or 5% dextrose, ZERBAXA is stable for 24 hours when stored at room temperature or 7 days when stored under refrigeration at 2°C to 8°C (36°F to 46°F). Discard unused portion.



Constituted ZERBAXA solution or diluted ZERBAXA infusion should not be frozen.

Important Safety Information (continued)

- Patients with renal impairment: Decreased efficacy of ZERBAXA has been observed in patients with baseline CrCl of 30 to ≤50 mL/min. In a clinical trial of adult patients, patients with cIAIs with CrCl >50 mL/min had a clinical cure rate of 85.2% when treated with ZERBAXA plus metronidazole vs 87.9% when treated with meropenem. In the same trial, patients with CrCl 30 to \leq 50 mL/min had a clinical cure rate of 47.8% when treated with ZERBAXA plus metronidazole vs 69.2% when treated with meropenem. A similar trend was also seen in the cUTI trial. Dose adjustment is required for adult patients with CrCl 50 mL/min or less. All doses of ZERBAXA are administered over 1 hour. Monitor CrCl at least daily in patients with changing renal function and adjust the dose of ZERBAXA accordingly.
- Hypersensitivity: ZERBAXA is contraindicated in patients with known serious hypersensitivity to the components of ZERBAXA (ceftolozane/ tazobactam), piperacillin/tazobactam, or other members of the beta-lactam class. Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving beta-lactam antibacterials. Before initiating therapy with ZERBAXA, make careful inquiry about previous hypersensitivity reactions to cephalosporins, penicillins, or other beta-lactams. If an anaphylactic reaction to ZERBAXA occurs, discontinue use and institute appropriate therapy.

Dosage and Administration

PREPARATION **OF SOLUTIONS**



How ZERBAXA **Is Supplied**

How ZERBAXA[®] (ceftolozane and tazobactam) is Supplied

ZERBAXA 1.5 g for injection is supplied in single-dose vials containing 1 g ceftolozane (equivalent to 1.147 g of ceftolozane sulfate) and 0.5 g tazobactam (equivalent to 0.537 g of tazobactam sodium) per vial, in a carton containing 10 vials.

Storage and Handling

ZERBAXA vials should be stored refrigerated at 2°C to 8°C (36°F to 46°F) and protected from light.

Package Size and Dimensions

Item Weight (Ounces per Vial)	Sales Package Size (Carton)	Carton Dimensions (L x H x W)
13.2 oz	Ten (10) single-dose vials	6 5/16" x 2 11/16" x 2 9/16"

Ordering

NDC for ZERBAXA IV Vials (10-ct)	67919-030-01
----------------------------------	--------------

NDC, National Drug Code.

ZERBAXA is available for purchase by all Merck Authorized Distributors, including, but not limited to, the distributors listed below. Contact your supplier for availability.

Wholesaler	Order Entry Number
Cencora (formerly known as AmerisourceBergen Corporation)	10237721
Besse Medical	10240631
Cardinal Health	5045810
DMS Pharmaceutical Group, Inc.	583420
FFF Enterprises	ZER003001
Henry Schein	5580056
McKesson Corporation	3971686
McKesson Medical Surgical	1015050
Metro Medical	661074
Morris & Dickson Co.	031179
Smith Drug Company	737924

Merck does not recommend the use of one authorized distributor over another. Merck does not make any warranty as to the services offered by any particular authorized distributor.

Important Safety Information (continued)

• Clostridioides difficile-associated diarrhea (CDAD), ranging from mild diarrhea to fatal colitis, has been reported with nearly all systemic antibacterial agents, including ZERBAXA. Careful medical history is necessary because CDAD has been reported to occur more than 2 months after the administration of antibacterial agents. If CDAD is confirmed, antibacterial use not directed against C. difficile should be discontinued, if possible.











Dosage and Administration

Preparation of Solutions

HOW ZERBAXA **IS SUPPLIED**

Coding and Billing Information

This resource contains a list of possible codes that may be relevant when billing for ZERBAXA[®] (ceftolozane and tazobactam). Please consult with the applicable payer or, where applicable, the Medicare Administrative Contractor, to understand the payer's specific billing requirements.

You are solely responsible for determining the appropriate codes and for any action you take in billing. Information about HCPCS codes is based on guidance issued by the Centers for Medicare & Medicaid Services (CMS) applicable to Medicare and may not apply to other public or private payers. Consult the relevant manual and/or other guidelines for a description of each code to determine the appropriateness for a particular code and for information on additional codes.

The information available here is compiled from sources believed to be accurate, but Merck makes no representation that it is accurate. This information is subject to change. Payer coding requirements may vary or change over time, so it is important to regularly check with each payer as to payer-specific requirements. The information available here is not intended to be conclusive or exhaustive, and is not intended to replace the guidance of a qualified professional advisor. Merck and its agents make no warranties or guarantees, expressed or implied, concerning the accuracy or appropriateness of this information for your particular use given the frequent changes in public and private payer billing. The use of this information does not guarantee payment or that any payment received will cover your costs.

HCPCS Codes for ZERBAXA¹

HCPCS Code	Description	Billing Units
J0695	Injection, ceftolozane 50 mg and tazobactam 25 mg	Each billing unit of ZERBAXA equal 50 mg of ceftolozane and 25 mg of
S9502	Home infusion therapy, antibiotic, antiviral, or antifungal therapy; once every 8 hours, administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem	N/A

HCPCS, Healthcare Common Procedure Coding System; APC, ambulatory payment classifications.

Note: Paid under Outpatient Prospective Payment System (OPPS) as a separate APC payment.²

Use form CMS-1500 for in-office injection; in the hospital outpatient department setting, submit claims using form UB-04 (also known as CMS 1450).

For questions on billing if a portion of the package is wasted, consult the applicable payer's policy regarding wastage. Please note that effective January 1, 2017, providers are required to use the JW modifier for Medicare claims with unused drugs or biologicals from single-use vials or single-use packages that are appropriately discarded.³

Important Safety Information (continued)

- Development of drug-resistant bacteria: Prescribing ZERBAXA in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.
- Adverse reactions in adult patients with HABP/VABP: The most common adverse reactions occurring in \geq 5% of adult patients receiving ZERBAXA in the HABP/VABP trial were hepatic transaminase increased (11.9%), renal impairment/renal failure (8.9%), and diarrhea (6.4%).
- Adverse reactions in adult patients with cIAI or cUTI: The most common adverse reactions occurring in ≥5% of adult patients receiving ZERBAXA in the cUTI and cIAI trials were headache (5.8%) in the cUTI trial, and nausea (7.9%), diarrhea (6.2%), and pyrexia (5.6%) in the cIAI trial.

IS of tazobactam



Dosage and Administration

Preparation of Solutions

How ZERBAXA **Is Supplied**

CODING AND BILLING INFORMATION

Coding and Billing Information (continued)

Drug Administration CPT[®] Code⁴

CPT Code	Description
96365	Intravenous infusion, for therapy, prophylaxis, or di or drug); initial, up to 1 hour

CPT, Current Procedural Terminology.

CPT © 2024 American Medical Association. All rights reserved. CPT is a registered trademark of the American Medical Association.

Possible Revenue Codes for Use in the Hospital Inpatient and Hospital Outpatient Settings⁵

Consult the relevant manual and/or other guidelines for a description of each code to determine the appropriateness of a particular code and for information on additional codes.

Revenue Code	Description
0636	Drugs requiring detailed coding
0250	General pharmacy

Important Safety Information (continued)

- Adverse reactions in pediatric patients with cIAI or cUTI: The most common adverse reactions occurring in $\geq 7\%$ of pediatric patients receiving ZERBAXA in the cIAI trial were diarrhea (17%), thrombocytosis (16%), pyrexia (13%), abdominal pain (11%), vomiting (10%), increased aspartate aminotransferase (7%), and anemia (7%). The most common adverse reactions occurring in \geq 7% of pediatric patients receiving ZERBAXA in the cUTI trial were thrombocytosis (9%), leukopenia (8%), diarrhea (7%), and pyrexia (7%).
- Pediatric Use: There is insufficient information to recommend dosage adjustment for pediatric patients younger than 18 years of age with cIAI and cUTI with eGFR 50 mL/min/1.73m² or less. ZERBAXA is not recommended in pediatric patients who have an eGFR 50 mL/min/1.73m² or less. Pediatric patients born at term or pre-term may not have an eGFR of 50 mL/min/1.73m² or greater at birth or within the first few months of life.

References:

1. Centers for Medicare & Medicaid Services. HCPCS Quarterly Update, July 2024. Accessed June 28, 2024. https://www.cms.gov/Medicare/ Coding/HCPCSReleaseCodeSets/HCPCS-Quarterly-Update 2. Centers for Medicare & Medicaid Services. Medicare Outpatient Hospitals Dataset: A Methodological Overview. May 2024. Accessed June 28, 2024. https://data.cms.gov/resources/medicare-outpatient-hospitals-methodology 3. Centers for Medicare & Medicaid Services. MLN Matters 13056. Accessed June 28, 2024. https://www.cms.gov/files/document/mm13056new-jz-claims-modifier-certain-medicare-part-b-drugs.pdf 4. American Academy of Professional Coders. CPT Code/Relative Value Search – 96365. Accessed June 28, 2024. https://www.aapc.com/codes/cpt-codes/96365 5. Noridian Healthcare Solutions. Revenue codes. Last updated March 18, 2024. Accessed June 28, 2024. https://med.noridianmedicare.com/web/jea/topics/claim-submission/revenue-codes



Copyright © 2024 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved. US-ZER-01883 07/24

agnosis (specify substance



Dosage and Administration

Preparation of Solutions

How ZERBAXA **Is Supplied**

CODING AND BILLING INFORMATION